

*Experiments*  
Учебен експеримент в природните науки

## SPECTRAL, MAGNETOCHEMICAL AND THERMOGRAVIMETRIC ANALYSIS (TGA) OF TRANSITION METAL COMPLEXES WITH N<sup>o</sup>O TYPES LIGANDS

G. Kumar, A. Munyaneza, G. Ditsie  
*University of Botswana, Botswana*

**Abstract.** Transition metal complexes, Na<sub>2</sub>[M(L<sub>1</sub>L<sub>2</sub>)]·H<sub>2</sub>O, where M = Cu(II), Co(II) and Ni(II) ions ; L<sub>1</sub> and L<sub>2</sub> are the anions of phenylalanine (Phe<sup>-</sup>) and nitrilotriacetic acid (NTA<sup>3-</sup>) respectively have been synthesized and characterized by elemental analysis, spectral (IR,UV in solidstate), magnetic susceptibility and thermogravimetric analysis (TGA). All the complexes have been suggested to show six fold octahedral geometry around the metal ion. Phenylalanine and nitrilotriacetic acid demonstrate bidentate and tetradentate behavior respectively, coordinating through their respective nitrogen atom and the oxygen atom (s) of the carboxylate group. Thermal behavior of Ni(II) complex is quite different from those of Cu(II) and Co(II) complexes which show similar type of thermal behavior. Based on their decomposition temperature, the thermal stability of the three complexes can be rated as Cu > Ni > Co.


**Keywords:** transition metal complexes, octahedral, phenylalanine, nitrilotriacetic acid, thermal analysis

### Introduction

Transition metal ions play a vital role in several biological processes in human body (Raja Balan et al., 2013). Divalent metals (Cu, Co and Ni) amino acid complexes have been of importance as model for the metal ligand systems and proved to be useful antibacterial agents against staphylococcus aureus, Escherichia coli, nutritive supplies for human and animals (Fitzsimmons et al., 1985; Martin et al., 1973; Rusu et al., 2009). Such complexes have also been found to act as active catalysts in many environmental and chemical reactions (Iakovidis et al., 1989). Studies on the synthesis, characterization and biological activities of mixed ligand transition metals complexes involving various amino acids including phenylalanine have been carried out by many researchers (Rusu et al., 2009; Sanap & Patil., 2013; Patil et al., 2012; Kabbani et al., 2005; Gupta & Srivastava., 1985; Reddy & Reddy., 2000; 2002; Shivankaret al., 2007; Kumar et al., 2013; 2014). Synthesis, spectral and magneto chemical studies of mixed

ligand amino acid chelates of divalent transition metals with nitrilotriacetic acid and glycine,  $\alpha$ -alanine, valine, or leucine have also been reported (Saxena & Srivastava, 1990). Studies on the octahedral structure of Cu(II), Co(II), Cd(II) and Zn(II) ternary complexes of nitrilotriacetic acid and alanine or phenylalanine prepared in acidic medium between pH 2.8 and 4.0 have been reported (Khalil et al., 2010). The formation, determination of structure, and the mode of coordination involved in a transition metal complex depend upon many factors such as, metal ions and their oxidation state, reaction medium and the structure of the ligands. Thus the, present paper describes the synthesis, spectral, magnetochemical and thermogravimetric studies of the mixed ligand Cu(II), Co(II) and Ni(II) complexes formed with nitrilotriacetic acid and phenylalanine between pH 4.20 and 8.93.

Structures of phenylalanine and nitrilotriacetic acid ligands are shown below in Fig. 1.



**Fig. 1.** Structure of the ligands

### Experimental

The two ligands, L-phenylalanine (Aldrich) and nitrilotriacetic acid (Fluka) and the metal salts,  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  (BDH),  $\text{CoCl}_2$  (Rochelle chemicals) and  $\text{NiCl}_2$  (Rochelle chemicals) were used as purchased without any further purification. Ligands were dissolved in one equivalent of sodium hydroxide (UNIAR) and metal ions solution were prepared in one equivalent of hydrochloric acid (Rochelle chemicals). IR spectra of ligands and synthesized metal complexes were recorded on a Perkin – Elmer FT-IR 2000 spectrophotometer in  $4000 - 400 \text{ cm}^{-1}$  range using KBr disc. Vario El C H N O/S elemental analyzer was used for the elemental analysis (C H N) of the complexes and magnetic susceptibility measurements were carried out on Johnson Matthey Alfa product magnetic susceptibility balance (University of Botswana). Electronic spectra of the complexes in solid state were recorded in the range 200-1300 nm (UV-NIR) on a

Perkin-Elmer UV/Vis/NIR spectrometer Lambda 750 and thermogravimetric analysis (TGA) was carried out on Mettler Toledo TGA/SDTA 851 instrument between 20° C and 1000°C temperature at a heating rate of 10° C /minute under the nitrogen atmosphere (Indian Institute of Sciences, Bangalore, India).

#### *Synthesis of complexes*

Equimolar (0.1M) solutions of ligands were mixed with metal ion (0.1M) solution separately in 1:1:1 molar ratio and the pH of the mixture solution was adjusted to an appropriate value between 4.20 and 8.93 to synthesize Cu(II), Co(II) and Ni(II) complexes.

#### *Copper(II) complex*

After mixing the above solutions, the pH of the resulting mixture solution was adjusted to ~ 4.20 by adding sodium hydroxide solution. The color of the solution remained blue throughout the experiment. The dark blue solution was then concentrated on a steam bath to reduce the volume of the solution between 25-30 ml and allowed to crystallize overnight. Blue crystalline product was then filtered and washed first with 50% alcohol-water mixture followed by acetone and dried in a vacuum desiccator.

#### *Cobalt(II) complex*

After mixing the solutions of both ligands and CoCl<sub>2</sub> solution, the pH of the resulting mixture solution was adjusted to ~ 8.93 by the addition of sodium hydroxide. The pink solution was then concentrated on a steam bath to reduce the volume between 20-25 ml and allowed to crystallize for about 24 hours. The pink powder product was then filtered and washed first with 50% alcohol-water mixture followed by acetone and dried in a vacuum desiccator.

#### *Nickel(II) complex*

The pH of the mixed solution of both ligands and NiCl<sub>2</sub> was adjusted to ~ 6.93 by adding sodium hydroxide. Bright blue solution was then concentrated on a steam bath and the volume of the solution was reduced between 25-30 ml and allowed to crystallize for about 24 hours. Light green powder product was then filtered and washed first with 50% alcohol – water mixture followed by acetone and dried in a vacuum desiccator.

## **Results and discussion**

### *Elemental analysis*

Elemental analysis (C H N) analytical data Calcd (found) for the suggested formula, Na<sub>2</sub>[M{C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CH(NH<sub>2</sub>)COO}{N(CH<sub>2</sub>COO)<sub>3</sub>}]H<sub>2</sub>O, [M=Cu(II), Co(II), Ni(II)], of the metal complexes and their respective color are given in Table 1.

**Table 1.** Elemental analysis and magnetic measurements data for the metal complexes

| Complex                                                                                                                                 | Color       | Calcd(found)%    |                |                | Magnetic measurements |                          |                          |                          |                    |
|-----------------------------------------------------------------------------------------------------------------------------------------|-------------|------------------|----------------|----------------|-----------------------|--------------------------|--------------------------|--------------------------|--------------------|
|                                                                                                                                         |             | C                | H              | N              | Temp (K)              | $\chi_g \times 10^6$ cgs | $\chi_M \times 10^6$ cgs | $\chi_M' \times 10^6$ BM | $\mu_{\text{eff}}$ |
| Na <sub>2</sub> [Cu(Phe)(NTA)].H <sub>2</sub> O<br>Na <sub>2</sub> [Cu(C <sub>15</sub> H <sub>18</sub> N <sub>2</sub> O <sub>9</sub> )] | Blue        | 37.54<br>(37.55) | 3.79<br>(3.49) | 5.83<br>(5.56) | 302                   | 2.357                    | 1131.19                  | 1252.15                  | 1.73               |
| Na <sub>2</sub> [Co(Phe)(NTA)].H <sub>2</sub> O<br>Na <sub>2</sub> [Co(C <sub>15</sub> H <sub>18</sub> N <sub>2</sub> O <sub>9</sub> )] | Pink        | 37.91<br>(38.42) | 3.82<br>(3.43) | 5.89<br>(5.90) | 301                   | 18.63                    | 8858.18                  | 8979.14                  | 4.65               |
| Na <sub>2</sub> [Ni(Phe)(NTA)].H <sub>2</sub> O<br>Na <sub>2</sub> [Ni(C <sub>15</sub> H <sub>18</sub> N <sub>2</sub> O <sub>9</sub> )] | Light green | 37.92<br>(37.80) | 3.83<br>(4.21) | 5.89<br>(5.69) | 297                   | 1012.77                  | 4811.37                  | 4932.33                  | 3.42               |

**Note:** Phe and NTA represent anions of phenylalanine and nitrilotriacetic acid respectively.

#### *Infrared studies*

In an earlier report (Khalil et al., 2010), the presence of bands in 1738-1728 cm<sup>-1</sup> range in the IR spectra of ternary Co(II), Ni(II) and Cu(II) complexes was attributed to the presence of uncoordinated free carboxylic acid group of nitrilotriacetic acid and hence it has been shown to act as a tridentate ligand in the octahedral complexes coordinating through the oxygen of two carboxylic acid groups and the nitrogen atom. Phenylalanine has been suggested to act as a monodentate ligand.

In the present IR studies, nitrilotriacetic acid shows characteristic  $\nu(\text{OH})$  absorption band for COOH group at ~1712 cm<sup>-1</sup> which vanishes in case of metal complexes. Instead,  $\nu_{\text{as}}(\text{COO}^-)$  and  $\nu_{\text{s}}(\text{COO}^-)$  absorption bands are observed between 1614-1582 cm<sup>-1</sup> and 1400-1383 cm<sup>-1</sup> range respectively (Saxena & Srivastava., 1990; Tomita & Uneo., 1963). The absence of bands in 1738-1728 cm<sup>-1</sup> range in the IR spectra of the metal complexes does not show the possibility of the presence of any uncoordinated carboxylic acid group of the nitrilotriacetic acid. In nitrilotriacetic acid,  $\nu(\text{CN})$  band is observed at 1331 cm<sup>-1</sup> which lowers in case of metal complexes. Hence, nitrilotriacetic acid coordinates through the nitrogen and oxygen atom of the carboxylate groups with the metal ions. In the free phenylalanine, the asymmetric (N-H) and symmetric (N-H) bands observed at ~3040 cm<sup>-1</sup> and ~2963 cm<sup>-1</sup> respectively are shifted to higher frequencies in the metal complexes indicates that the nitrogen of the amino group coor-

dinates with the metal ions (Shivankar et al., 2007). The IR spectrum of phenylalanine also shows  $\nu_{as}(\text{COO}^-)$  and  $\nu_s(\text{COO}^-)$  bands at  $1556\text{ cm}^{-1}$  and  $1408\text{ cm}^{-1}$  respectively. In metal complexes, the shifting of  $\nu_{as}(\text{COO}^-)$  band to the higher frequency range ( $1614\text{--}1582\text{ cm}^{-1}$ ) and  $\nu_s(\text{COO}^-)$  band to the lower frequency range ( $1400\text{--}1383\text{ cm}^{-1}$ ) suggest the coordination of the carboxylic group via oxygen atom. In addition, metal complexes also show characteristic  $\nu(\text{OH})$  frequency around  $\sim 3300\text{ cm}^{-1}$  in the region  $3400\text{--}3200\text{ cm}^{-1}$  attributed to the lattice water molecule and a strong band in the region  $1614\text{--}1550\text{ cm}^{-1}$  is also the indicative of the presence of water molecule (Shivankar et al., 2007; Bellamy, 1975; Nakamoto, 1970). The difference in the  $\text{COO}^-$  frequencies for the complexes are greater than the difference in the corresponding frequencies in phenylalanine suggesting monodentate behavior of its carboxylic acid group (Devereux et al., 1998; Mitie et al., 2009; Danyi et al., 2006). Thus it can be inferred that the coordination in metal complexes takes place through the oxygen atom of the carboxylic acid group and the nitrogen of the amino group, suggesting a bidentate behavior for the phenylalanine. However, tentative assignment of the main IR bands for the ligands and metal complexes are given in Table 2.

**Table 2.** Main IR frequencies ( $\text{cm}^{-1}$ ) of ligands and their metal complexes


| Compound                                                                                                                                              | $\nu(\text{OH})$ | $\nu_{as}$<br>(N-H) | $\nu_{as}$<br>(N-H) | $\nu_{as}$<br>$\text{COO}^-$ | $\nu_s$ $\text{COO}^-$ | $\nu$ CO<br>$\text{COOH}$ | $\nu$ HOH | $\nu_s$ (CH) | $\nu_s$<br>(CN) |
|-------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|---------------------|---------------------|------------------------------|------------------------|---------------------------|-----------|--------------|-----------------|
| Phenylalanine                                                                                                                                         | -                | 3040                | 2963                | 1556                         | 1408                   | -                         | -         | -            | -               |
| Nitrilotriacetic acid                                                                                                                                 | -                | -                   | -                   | -                            | -                      | 1712                      | -         | 3040<br>2992 | 1331            |
| $\text{Na}_2[\text{Cu}(\text{Phe})(\text{NTA})] \cdot \text{H}_2\text{O}$<br>$\text{Na}_2[\text{Cu}(\text{C}_{15}\text{H}_{18}\text{N}_2\text{O}_9)]$ | 3247             | 3173                | 3032                | 1614                         | 1383                   | -                         | 1614      | 2962         | 1322            |
| $\text{Na}_2[\text{Co}(\text{Phe})(\text{NTA})] \cdot \text{H}_2\text{O}$<br>$\text{Na}_2[\text{Co}(\text{C}_{15}\text{H}_{18}\text{N}_2\text{O}_9)]$ | 3358             | 3172                | 2967                | 1579                         | 1396                   | -                         | 1578      | 2967         | 1323            |
| $\text{Na}_2[\text{Ni}(\text{Phe})(\text{NTA})] \cdot \text{H}_2\text{O}$<br>$\text{Na}_2[\text{Ni}(\text{C}_{15}\text{H}_{18}\text{N}_2\text{O}_9)]$ | 3354             | 3291                | 3022                | 1582                         | 1400                   | -                         | 1582      | 2942         | 1322            |

**Note:** Phe and NTA represent anions of phenylalanine and nitrilotriacetic acid respectively.

## Magnetic measurements and electronic spectra

### *Copper(II) complex*

The observed magnetic moment for the copper(II) complex is 1.73 B.M which is well in the expected range (1.70 – 2.20 BM) for the presence of one unpaired electron corresponds to the spin free octahedral geometry for the complex (Figgis & Lewis, 1960). Copper(II) with  $d^9$  configuration shows main absorption band for the d-d transition at 657 nm ( $15\,220\text{ cm}^{-1}$ ) which can safely be assigned to  ${}^2E_g \rightarrow {}^2T_{2g}$  transition and a charge transfer band is observed at 265 nm ( $37\,735\text{ cm}^{-1}$ ) (Lever, 1984). The electronic spectra of copper(II) complex is shown in Fig. 2.




**Fig. 2.** Electronic spectrum of copper(II) complex

### *Cobalt(II) complex*

The observed magnetic moment, 4.65 BM for cobalt(II) complex suggests a six fold octahedral structure (Figgis & Lewis, 1960; Cotton et al., 1999). This value is higher than the required value for a free spin  $d^7$  state corresponding to three unpaired electrons. Co(II) ion with  $d^7$  configuration with an octahedral geometry should show three absorption bands corresponding to three spin allowed transitions in its electronic spectrum; ( $\nu_1$ )  ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$  observable near infrared region; ( $\nu_2$ )  ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$  and ( $\nu_3$ )  ${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$ . Since,  $\nu_2$  transition involves a two electron process and should be

weaker by  $\sim 10^{-2}$  than the other transitions (Cotton et al., 1999), hence its position may however be ambiguous and is rather difficult to locate with certainty (Lever, 1984; Cotton et al., 1999). Sometimes it appears as a shoulder on either side of the main absorption band ( $\nu_3$  transition) and in some cases it may be unobservable. Besides, the spectra should also show charge transfer bands. In the present cobalt (II) complex, the main absorption band ( $\nu_3$ ) is observed at 509 nm ( $19\,646\text{ cm}^{-1}$ ) which can safely be assigned to  ${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$  transition. The band observed at 1188 nm ( $8418\text{ cm}^{-1}$ ) in the NIR can be attributed to ( $\nu_1$ ),  ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$  transition. The charge transfer band is observed at 261 nm ( $38\,314\text{ cm}^{-1}$ ). The electronic spectra of cobalt (II) complex is shown in Fig. 3.




**Fig. 3.** Electronic spectrum of cobalt(II) complex

#### *Nickel(II) complex*

The observed magnetic moment of 3.42 BM indicates an octahedral structure ( $d^8$  configuration) for the nickel(II) complex (Figgis & Lewis., 1960; Cotton et al., 1999), which is supported by its electronic spectrum. Ni(II) has a  $d^8$  electronic configuration and therefore three d-d electronic transitions are expected to occur in addition to charge transfer bands:  ${}^3A_{2g} \rightarrow {}^3T_{1g}(P)$ ;  ${}^3A_{2g} \rightarrow {}^3T_{1g}(F)$  and  ${}^3A_2 \rightarrow {}^3T_{2g}(F)$ . The spectrum

for Ni(II) complex is presented in Fig. 4. The maximum at 627 nm ( $15\,948\text{ cm}^{-1}$ ) is assigned to  ${}^3\text{A}_{2g} \rightarrow {}^3\text{T}_{1g}(\text{F})$  transition and the one at 388 nm ( $25\,773\text{ cm}^{-1}$ ) corresponds to  ${}^3\text{A}_{2g} \rightarrow {}^3\text{T}_{2g}(\text{P})$  transition.  ${}^3\text{A}_{2g} \rightarrow {}^3\text{T}_{2g}$  transition is observed at 1070 nm ( $9345\text{ cm}^{-1}$ ). Charge transfer bands are seen at 258 nm ( $38\,759\text{ cm}^{-1}$ ) (Cotton et al., 1999).



**Fig.4.** Electronic spectrum of nickel(II) complex


Magnetic susceptibility measurements together with elemental analysis (C,H,N) of the metal complexes are given in Table1.

Based on the IR results as well as on the magnetic measurements, elemental analysis and electronic spectra, we suggest the general structure of the synthesized complexes as shown in Fig. 5.

### Thermogravimetric analysis (TGA) study


Thermogravimetric analysis is used to determine the thermal stability and the fragmentation patterns of materials (Vega-Lizama et al., 2015; Hu et al., 2015). The thermal behavior of copper, cobalt and nickel complexes with phenylalanine and nitrilotriacetic

acid was studied in the temperature range 20-1000°C. As a general observation, the fragmentation for all the complexes starts with the loss of lattice water molecule and ends with the metal carbonate residue,  $\text{MCO}_3$  ( $\text{M} = \text{Cu}, \text{Co}, \text{Ni}$ ) (Figs. 6-8). The copper complex showed a better thermal stability than the other two complexes ( $\sim 105^\circ\text{C}$ ) while cobalt complex was the least stable ( $\sim 80^\circ\text{C}$ ). The nickel complex was stable up to  $95^\circ\text{C}$ . Table 3 shows detailed fragmentation patterns for the three complexes.




**Fig. 5.** Structure of complexes; where  $\text{M} = \text{Co(II)}, \text{Ni(II)}, \text{Cu(II)}$  ions


The fragmentation pathways as shown by TGA study confirm the structure of the complexes as suggested in Fig. 5.



**Fig. 6.** TGA plot for Cu – Phe – NTA complex



**Fig. 7.** TGA plot for Co - Phe - NTA complex



**Fig. 8.** TGA plot for Ni - Phe - NTA complex

**Table 3.** Thermogravimetric analysis results for M – Phe - NTA

| Complex   | Stability (° C) | Temp. range (° C) | Removed group(s)*                                       | Mass loss (%) |            |
|-----------|-----------------|-------------------|---------------------------------------------------------|---------------|------------|
|           |                 |                   |                                                         | Found         | Calculated |
| Cu-Ph-NTA | 105             | 105-169           | H <sub>2</sub> O                                        | 3.82          | 3.75       |
|           |                 | 169-318           | Na <sub>2</sub> , Ph-CH <sub>2</sub> -CHNH <sub>2</sub> | 37.21         | 38.34      |
|           |                 | 318-412           | OOCHCH <sub>2</sub>                                     | 48.87         | 50.42      |
|           |                 | 412-625           | COO                                                     | 58.00         | 59.59      |
|           |                 | 914-              | Remaining residue, CuCO <sub>3</sub>                    | 24.68         | 25.62      |
| Co-Ph-NTA | 80              | 80-129            | H <sub>2</sub> O                                        | 3.82          | 3.79       |
|           |                 | 129-181           | Na                                                      | 7.61          | 8.63       |
|           |                 | 181-308           | Na, OOCCHCH <sub>2</sub>                                | 26.82         | 25.68      |
|           |                 | 308-436           | Ph-CH <sub>2</sub> -CHNH <sub>2</sub> , COO             | 59.87         | 60.21      |
|           |                 | 987-              | Remaining residue, CoCO <sub>3</sub>                    | 25.93         | 25.04      |
| Ni-Ph-NTA | 95              | 95-208            | H <sub>2</sub> O                                        | 3.18          | 3.79       |
|           |                 | 208-269           | Ph-CH <sub>2</sub>                                      | 22.73         | 22.96      |
|           |                 | 269-357           | Na <sub>2</sub> , OOCCH <sub>2</sub>                    | 45.08         | 44.87      |
|           |                 | 951-              | Remaining residue, NiCO <sub>3</sub>                    | 24.68         | 24.85      |

\*The groups removed from the second step onwards include the groups removed in the preceding steps.

### Conclusion

In view of the evidences obtained above from the IR, electronic spectra, magnetic susceptibility measurements and thermogravimetric analysis, a sixfold octahedral structure has been suggested for all the complexes in which nitrilotriacetic acid and phenylalanine act as a tetradentate and bidentate ligands respectively. Both the ligands coordinate with their respective nitrogen atom and the oxygen atom(s) of the carboxylic acid groups. Based on the temperature decomposition, the thermal stability of the complexes is suggested as Cu > Ni > Co. The suggested structure for the complexes is shown in Fig. 5.

**Acknowledgements:** Authors gratefully acknowledge the assistance provided by the University of Botswana, the Third World Academy of Science (TWAS) and the India Institute of Sciences, India.

### REFERENCES

- Bellamy, L.J. (1975). *The infra red spectra of complex molecules*. London: Chapman.
- Cotton, F.A., Wilkinson, G., Murillo, C.A. & Bochmann, M. (1999). *Advanced inorganic chemistry*. New York: Wiley.
- Danyi, W., Ning, L., Gui, L. & Kemin, Y. (2006). Synthesis, catalytic and biological activity of novel dinuclear copper complex with Schiff base. *Science in China B, Chemistry*, 49(3), 225-229.

- Devereux, M., Jackman, M., McCann, M. & Casey, M. (1998). Preparation and catalase-type activity of manganese(II) amino acid complexes. *Polyhedron*, 17, 153-158.
- Figgis, B.N. & Lewis, J. (1960). The magnetochemistry of complex compounds (pp 400-454). In: Lewis, J. & Wilkins, R. G. (Eds.). *Modern coordination chemistry*. New York: Interscience.
- Fitzsimmons, B.W., Hume, A., Larkworthy, L.F., Turnbull, M.H. & Yavari, A. (1985). The preparation and characterization of some complexes of iron(II) with amino acids. *Inorg. Chim. Acta*, 106, 109-114.
- Gupta, M. & Srivastava, M.N. (1985). Synthesis and characterization of mixed ligand complexes of copper(II), nickel(II), cobalt(II) and zinc(II) with glycine and uracil or 2-thiourecil. *Polyhedron*, 4, 475-479.
- Hu, S., Ma X., Lin Y., Yu, Z. & Fang S. (2015). Thermogravimetric analysis of the combustion of paper mill sludge and municipal solid waste. *Energy Conversion & Management*, 99, 112-118.
- Iakovidis, A., Hadjiliadis, N., Schollhorn, H., Thewalt, U & Trotscher, G. (1989). Interaction of cis-Pt(NH<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> with amino acids: the crystal structure of cis-[Pt(NH<sub>3</sub>)<sub>2</sub>(gly)](NO<sub>3</sub>), cis-[Pt(NH<sub>3</sub>)<sub>2</sub>(ala)](NO<sub>3</sub>), cis-[Pt(NH<sub>3</sub>)<sub>2</sub>(val)](NO<sub>3</sub>). *Inorg. Chim. Acta.*, 164, 221-229.
- Kabbani, A.T., Ramadan, H., Hammud, H.H., Ghannoum, A.M. & Mouneimne, Y. (2005). Synthesis of some metal complexes of N-[(benzoylamino)-thioxomethyl]-amino acid (HL). *J. Univ. Chem. Tech. & Metallurgy*, 40(4), 339-344.
- Khalil, M.M.H., Ismail, E.H., Azim, S.A. & Souaya, E.R. (2010). Synthesis, characterization and thermal analysis of ternary complexes of nitrilotriacetic acid and alanine or phenylalanine with some transition metals. *J. Therm. Anal. Calorim.*, 101, 129-135.
- Kumar, G., Munyaneza, A. & Ditsie, G. (2013). Synthesis and characterization of mixed ligand Cu(II) and Co(II) complexes of L-phenylalanine and iminodiacetic acid. *Chemistry*, 22, 556-563.
- Kumar, G., Munyaneza, A. & Ditsie, G. (2014). Synthesis and characterization of mixed ligand Co(II) and Ni(II) complexes of L-phenylalanine and hippuric acid. *Chemistry*, 23, 881-889.
- Lever, A.B.P. (1984). *Inorganic electronic spectroscopy*. New York: Elsevier.
- Martin, R.P., Petit-Ramel, M.M. & Scharff, J.P. (1973). Mixed-ligand metal ion complexes of amino acids and peptides (pp. 1- 2). In: Sigel, H. (Ed.). *Metals in biological systems; vol. 2: mixed ligand complexes*. New York: Marcel Dekker.
- Mitie, D., Milenkovic, M., Milosavelevic, S., Godevac, D., Miodragovic, Z., Andelkovic, K. & Miodragovic, D. (2009). Synthesis, characterization and antimicrobial activity of Co(II), Zn(II) and Cd(II) complexes with N-benzoyloxy carbonyl-S-phenylalanine. *Eur. J. Med. Chem.*, 44, 1537-1544.

- Nakamoto, K. (1970). *Infrared spectra of inorganic and coordination compounds*. New York: Wiley.
- Patil, A.R., Donde, K.J., Raout, S.S., Patil, V R. & Lokhande, R.S. (2012). Synthesis, characterization and biological activity of mixed ligand Co(II) complexes of Schiff base 2-amino-4-nitrophenol-n-salicylidene with some amino acids. *J. Chem. Pharma Res.* 4, 1413-1425.
- Raja Balan, A.M.K., Ashok, R.F.N., Vasanthi, M., Prabu, R. & Pushpraj, A. (2013). Mixed ligand complexes of nickel(II), copper(II) and zinc(II) with nicotinalide and thio-cynate. *Intern. J. Life Sci. & Pharma Res.*, 3(2), L-67-L-75.
- Reddy, P.R. & Reddy, A.M. (2000). Synthesis and characterization of mixed ligand complexes of bio metals with pyrimidine nucleoside (uridine) and amino acids. *Proc. Indian. Acad. Sci. (Chem. Sci.)*, 112(6), 593 - 600.
- Reddy, P.R. & Reddy, A.M. (2002). Synthesis and characterization of mixed ligand complexes of Cu(II), Ni(II) and Co(II) with cytidine and amino acids. *Indian. J. Chem.*, 41A, 2083-2087.
- Rusu, D., Stanila, A., Marian, I.O., Marian, C.O., Rusu, M. & Lucaciu, R. (2009) Synthesis and characterization of some cobalt(II) complexes with amino acids having biological activities. *Rev. Chim. (Bucuresti)*, 60, 939-943.
- Sanap, S.V. & Patil, R.M. (2013). Synthesis, characterization and biological activity of chiral mixed ligand Ni(II) complexes. *Res. J. Pharma. Sci.* 2(1), 1-10.
- Saxena, V.K. & Srivastava, M.N. (1990). PMR spectral studies of mixed ligand amino acid chelates of cobalt(II), nickel(II), copper(II) and zinc(II) with nitrilotriacetic acid and glycine,  $\alpha$ -alanine, valine, or leucine. *J. Inorg. Bio-Chem.*, 38, 37-46.
- Shivankar, V.S., Vaidya, R.B., Dharwadkar, S.R. & Thakkar, N.V. (2007). Synthesis, characterization and biological activity of mixed ligand Co(II) complexes of 8-hydroxy-quinoline and some amino acids. *Synth. React. Inorg. Met-Org. Chem.*, 33(9), 1597-1622.
- Tomita, Y. & Uneo, K. (1963). The properties and infrared spectra of nitrilotriacetate chelates. *Bull. Chem. Soc. Japan.* 36, 1069 – 1073.
- Vega-Lizama, T., Diaz-Ballote, L., Hernandez-Mezquita, E., May-Crespo, F., Castro-Borges, P., Castillo-Atoche, A., Garcia-Gonzales, G. & Maldonado, L. (2015). Thermogravimetric analysis as a rapid and simple method to determine the degradation degree of soy biodiesel. *Fuel*, 156, 158-162.

✉ **Dr. G. Kumar (corresponding author)**

Department of Chemistry  
University of Botswana  
P/Bag UB 704, Gaborone, Botswana  
E-mail: kumar@mopipi.ub.bw